

Arrival Time

AM Bowel Preparation with SuTab

- **You must be accompanied by a friend or relative to drive and assist you home. We WILL NOT discharge you to a cab, bus, uber or other transportation service without a responsible party with you. You may not drive until the day after your procedure.**

Special Instructions Regarding Medications

- If you are on an **anticoagulant/blood thinner** our nurse will contact you with instructions after contacting your prescribing physician. If you have not heard from the nurse 1 week prior to your procedure please contact our office.
- **Do Not take Iron supplements for one week prior** to the procedure date. Multivitamins are OK.
- **Diabetic patients:**
- **Do Not take Water Pill/Diuretic** (Lasix (furosemide), HCTZ (Hydrochlorothiazide), Dyazide, Diovan HCT, Spironolactone), **on the day of the procedure**
- **Heart/Blood Pressure/Thyroid/Seizure/Anxiety Medications can be taken on the day of the procedure** with a sip of water.
- **Stop Effexor one day before** the procedure (if applicable)
- **Asthmatic patients** should bring their **INHALERS** with them on the day of the procedure



THE DAY BEFORE YOUR PROCEDURE

IMPORTANT NOTE: follow the instructions below, NOT the instructions on the prep's packaging

- **DO NOT EAT ANY SOLID FOODS. YOU MAY ONLY HAVE CLEAR LIQUIDS**
- Take all of your normal prescription medications, **unless advised otherwise.**

- **At 6 pm** – Open 1 bottle of 12 SuTab tablets. Fill provided container with 16 oz of **Water**
Swallow each tablet with a sip of water and drink the entire amount over 15-20 minutes
- Approximately one (1) hour after the last tablet is ingested, fill the container with 16 oz of Water and drink the Entire amount over 30 minutes.
- Approximately 30 minutes after finishing the second container of water, fill the container again with 16 oz water And drink the entire amount over 30 minutes.

DAY OF YOUR PROCEDURE

- **At 12 am Midnight -**
– Open 1 bottle of 12 SuTab tablets. Fill provided container with 16 oz of **Water**
Swallow each tablet with a sip of water and drink the entire amount over 15-20 minutes
- Approximately one (1) hour after the last tablet is ingested, fill the container with 16 oz of Water and drink the Entire amount over 30 minutes.
- Approximately 30 minutes after finishing the second container of water, fill the container again with 16 oz water And drink the entire amount over 30 minutes.

NOTHING BY MOUTH 4 HOURS BEFORE YOUR ARRIVAL TIME

General or reschedule questions call 610-431-3122