



GI-ASC LLC
1311 BRISTOL Pike Ste 130
BENSALEM, PA 19020
T (215) 632-3500 ext. 5114

Colonoscopy Instructions

Miralax Prep

PATIENT NAME: _____ DOB: _____

PHYSICIAN: _____

PROCEDURE DATE: _____ PROCEDURE TIME: _____

PLEASE ARRIVE 1/2HR PRIOR TO YOUR PROCEDURE TIME FOR CHECK-IN

LOCATION: _____

OFFICE PHONE NUMBER 215-632-3500 ~ HOURS 8:00 AM – 5:00 PM

YOUR MEDICATION INSTRUCTIONS: _____ **initial here**

- Stop** taking medications with Iron 5 days prior to the procedure.
- Continue all prescribed medications unless otherwise instructed.
- The following medications may need to be stopped 5-7 days prior to your procedure.
Please discuss this with your prescribing physician & cardiologist: blood thinners such as: coumadin, aspirin, plavix, aggrenox, xarelto, eliquis, pradaxa, arthritis medications, motrin ibuprofen, naprosyn, celebrex etc.
- Stop any blood thinning medications only if you were instructed to do so.
- Take your medications 3 or more hours before your scheduled procedure with a sip of water. If you are not able to take them 3 hours before your procedure, please wait until after your procedure.
- Do not drink any fluid within 4 hours of your procedure or it may be rescheduled.**
Sips of water for medications mentioned above are allowed.
- Diabetics:** If you have diabetes, check with your physician to see if changes are needed.
Do not take your usual oral diabetic medications the morning of your procedure. Check your blood sugar before arrival. Call the office 215-632-3500 if your blood sugar is 70 or below. Notify the staff immediately upon arrival as well.

SPECIAL MEDICATION INSTRUCTIONS:

What is a Colonoscopy?

A colonoscopy is a test that allows the physician to examine the lining of the large intestine with a thin flexible instrument. Polyps, or small growths, can be removed to prevent colon cancer. Biopsies can be taken to diagnose other diseases.

- *An intravenous (IV) catheter will be inserted into a vein in your arm to provide hydration and medication to keep you asleep during your procedure.*
- *You will receive nasal oxygen. We will monitor your heart rate, blood pressure and blood oxygen level throughout the procedure.*
- *Air or carbon dioxide and water will be placed into your colon during the exam. It is normal to expel these after the procedure.*
- *You will rest in the recovery room after the procedure until you can tolerate liquids, walk unassisted, and have stable vital signs.*

THE ENTIRE PROCESS TAKES ABOUT 2 HOURS.

If the start of your procedure is delayed, we will do our best to keep you informed. Rest assured that during your colonoscopy you will receive the highest quality care.

IMPORTANT

72 HOURS NOTICE IS REQUIRED FOR CANCELLATION.

IF YOU CANCEL LESS THAN 72 HOURS BEFORE YOUR PROCEDURE, YOU WILL BE CONSIDERED A “NO SHOW” AND MAY BE CHARGED A \$ 200.00 FEE. THIS FEE IS NOT COVERED BY INSURANCE.

PREPARATION: 1 Week before your procedure

- ❑ You will receive anesthesia during the procedure, *so a responsible adult MUST accompany you to your colonoscopy. This person MUST bring you to the office, stay in the building the entire time, then bring you home. The side effects of anesthesia are sometimes dizziness, forgetfulness, and sleepiness therefore:*
 - You may NOT be dropped off for your procedure.
 - A responsible adult must accompany you and stay with you if you take public transportation.
 - Bus, taxi, shuttle, or UBER drivers CANNOT be your responsible driver.
 - If you do not have a responsible adult to accompany and stay with you, your procedure will be rescheduled.

- ❑ **Medication: See page 1.**

- ❑ **Purchase Prep:**

For your convenience, a Colonoscopy Prep Kit can be purchased at the office. This Kit includes the necessary prep, Simethicone and 4 laxative tablets.
Please read directions carefully when mixing your prep.
For mixing instructions & prep timing, please see pages 5 & 6 of this packet

Or you may purchase your own:



Miralax Laxative Powder (2 (two) 238g or 8.3 oz bottles)

Clear juice – apple, white grape, lemonade

Gatorade or similar sports drinks

Kool Aid, Crystal Light

★ **NO RED OR PURPLE JUICE**

★ **NO JUICE WITH PULP**

★ **NO DAIRY DRINKS**

★ **NO ALCOHOL**

Dulcolax – 4 (four) 5 mg tablets

Simethicone (Gas-X) 2 (two) 125 mg tablets

PREPARATION: 2-3 Days before your procedure

DO:

Drink plenty of fluids – at least eight 8 oz glasses per day.

This will help avoid dehydration, make the prep work more effectively and make you feel better overall.



DO NOT:

Do not eat any of the following foods, as they can remain in your colon after the completion of your prep:

- Seeds
- Nuts
- Oatmeal
- Whole grains
- Beans
- Peas
- Corn
- Peels of fruits or vegetables

A successful colonoscopy is a collaboration between you and your physician. Following these instructions is necessary so the physician can properly perform the procedure.

Failure to follow these directions may result in a poor prep and the need for your procedure to be repeated or rescheduled.

Medications:

Review any personalized instructions on page 1.

Responsible Adult:

Review requirement for the person accompanying you to your procedure on page two. **Remember, a responsible adult must accompany you to your colonoscopy and stay until discharge. Visitors may be asked to wait in their car due to social distancing requirements.**

Other Items:

- Wear comfortable clothes. You may feel bloated after the procedure.
- Leave valuables at home or with the adult accompanying you.
- Remove all jewelry, including body piercings.

PREPARATION: The day before your procedure

<p style="text-align: center;">The morning of your Colonoscopy</p> <p style="text-align: center;">NO SOLID FOOD</p>	<p>Follow a diet of <u>CLEAR LIQUIDS ONLY</u>:</p> <ul style="list-style-type: none"> * Water * Coffee or tea, <i>without milk or cream</i> * Clear juice – apple, white grape, lemonade (no pulp) * Broth, bouillon * Gatorade or similar sports drinks * Kool Aid, Crystal Light * Carbonated Drinks – 7-UP, GINGER ALE, SPRITE & ORANGE ARE ACCEPTABLE * Jell-O™ (no added fruit or toppings; not red or purple) * Popsicles (not red or purple) * NO RED OR PURPLE JUICE * NO JUICE WITH PULP * NO DAIRY DRINKS * NO ALCOHOL
<p style="text-align: center;">2:00 pm the Day Before your Colonoscopy</p> <p style="text-align: center;">NO SOLID FOOD</p>	<p>Take 4 (four) 5 mg Dulcolax Tablets with 8 oz. water.</p> <p><u>*Prepare the prep*</u>: Please follow these mixing instructions carefully: put 1 Gatorade packet, ½ of the Miralax bottle in the 64 oz jug & fill with water or clear liquid of your choice, shake to mix & refrigerate until you start the prepping process.</p> <p style="text-align: center;"><i>Find a nearby bathroom!</i></p>
<p style="text-align: center;">4:00-6:00 pm the Day Before your Colonoscopy</p> <p style="text-align: center;">NO SOLID FOOD</p>	<p>First Dose of Liquid Prep (128 oz):</p> <ul style="list-style-type: none"> ★ Drink 8 oz of liquid prep every 20 minutes until jug is completed. ★ Immediately following completion of above dose, repeat process as stated above to complete 2nd jug. ★ Clear liquid diet can be continued until your morning dose begins. ★ Try your best to do this to prevent dehydration.

After completion of 128 oz of liquid prep, prepare the remaining 64 oz per the mixing instructions above, using half the amount of Miralax, then refrigerate until morning dose. (You will have ½ bottle of Miralax remaining that will not be used)

PREPARATION: The day of your procedure

<p>6 HOURS BEFORE YOUR APPOINTMENT</p> <p>NO SOLID FOOD</p>	<p>Second Dose of Liquid Prep (64 oz):</p> <ul style="list-style-type: none">★ Drink 8 oz's of liquid prep every 15-30 minutes for 1-2 hours.★ Take 2 Simethicone (Gas-X) tablets after completing the Miralax drink.★ Try your best to do this to prevent dehydration.★ The goal is to have slightly yellow and liquid stool.
<p>4 HOURS BEFORE YOUR APPOINTMENT</p> <p>NO FOOD OR DRINK</p>	<p>All prep and clear liquids MUST be completed 4 hours prior to your appointment.</p> <p>This is the time when morning medications should be taken or wait until after your procedure.</p> <p>No additional liquids or your procedure may be rescheduled.</p> <p>No hard candy, chewing gum or medications.</p> <p>Nothing by mouth.</p>

AFTER YOUR COLONOSCOPY

For routine questions, call our office at 215-632-3500 between 8:00am and 5:00 pm.

IF YOU HAVE AN URGENT CONCERN, CALL 215-632-3500 TO REACH OUR ON-CALL PHYSICIAN AFTER HOURS.

- Rest and relax for the remainder of the day.
- We recommend that a responsible adult stay with you for 24 hours following your procedure.
- Resume your normal diet slowly and as tolerated.
- A feeling of fullness or cramping from remaining air or carbon dioxide may occur. This is totally normal. Mild activities such as walking will help to expel any excess gas. Lying on your left side or directly on your stomach may also help.
- Do not drive or operate any machinery, sign any legal documents, or make critical decisions or return to work until the next day.
- Do not drink alcohol or take any unprescribed medication.
- Mild abdominal discomfort or a small amount of rectal bleeding is not unusual after the procedure. However, if you experience significant pain, rectal bleeding, fever and vomiting or any other worrisome symptoms please notify your physician immediately.
- Follow up with your physician's recommendations for continued colon health.