

COLONOSCOPY – DIET AND LAXATIVE BOWEL PREPARATION – PEG3350

Hillmont GI. P.C. 215.402.0800

WHAT TO EXPECT:

- Most people have a bowel movement within an hour or two of starting the laxative. Sometimes, there may be a delay of several hours. Be patient and stay close to a bathroom. Multiple bowel movements will occur and may continue for several hours after you have finished drinking the solution
- Your stools should become clear of solid material. **When you are fully prepared the final bowel movements should look like urine in the toilet water**
- Bloating, chills and/or nausea are common after the first few glasses and more common if the liquids are chilled. This is usually temporary and improves once bowel movements begin. If you develop vomiting, severe discomfort or bloating, stop taking the prep for a while until the symptoms go away. If vomiting persists, stop the preparation and call our office 215-402-0800

7 DAYS PRIOR TO YOUR PROCEDURE: Review Medication handout, obtain laxative preparation

3 DAYS PRIOR TO YOUR PROCEDURE: AVOID WHOLE GRAINS, NUTS, CORN, POPCORN & SMALL SEEDS

- Whole grains- oatmeal, granola, 7 grain breads, rice, etc.
- Seeds - raspberries, strawberries, cucumbers, tomatoes, poppy, sesame, seeds, etc

THE DAY BEFORE THE PROCEDURE: NO SOLID FOOD AT ALL. CLEAR LIQUIDS ALL DAY

- NO CLOUDY LIQUIDS, milk, milk drinks, creamer, liquids that have pulp, NO alcoholic beverages

CLEAR LIQUIDS: include any liquid that you can see clearly through. Avoid RED and PURPLE

- | | |
|---------------------------------|---|
| • Water | • Coffee or Tea– without milk, non-dairy creamer or cream |
| • Carbonated beverages / soda | • Clear fruit juices without pulp – apple, grape, lemonade, etc |
| • Gatorade® Kool-aid® | • Clear hard candy, sugar, and lemon |
| • Jello® Popsicles or water ice | • Clear broth (chicken, beef, vegetable, etc) |

IMPORTANT NOTE: FOLLOW THESE INSTRUCTIONS AND NOT the instructions on the Prep’s packaging.

THE NIGHT PRIOR TO YOUR PROCEDURE: 7:00 pm

- Add water and flavor packets if available to the container and shake until medication is dissolved. Drink 8-ounces every 15 minutes until you have finished half of the container.
- Save the remainder of the preparation to drink later
- Drink at least an additional 16 ounces of clear liquid before bedtime. Drinking more usually leads to a better preparation and complete and thorough examination

THE DAY OF YOUR PROCEDURE: FIVE (5) hours before your procedure, AT _____ AM/PM

- Re-shake the remaining bowel prep until medication is dissolved. Drink 8 ounces every 15 minutes until the entire container is finished.
- Drink at least an additional 16- ounces of clear liquid before you have to stop drinking.
- You **MUST stop drinking 3 hours before** your procedure time, **so nothing after** _____
- NO SMOKING the day of the procedure
- **You must put NOTHING in your mouth for the last 3 hours prior to your procedure.** This means NO water, ice chips, gum, candy, cough drops, etc., **except that you should take your medications with a sip of water**

**For questions about the preparation, medications or to reschedule your appointment,
call 215-402-0800 between 9:00AM and 5:00PM**