US Digestive Health / Springfield ASC

1528 Bethlehem Pike, Flourtown, PA 19031 Phone: 215-402-0600

DATE:	TIME:
	PREPARATION FOR A FLEXIBLE SIGMOIDOSCOPY
For the	e best possible examination, the bowel must be empty of waste material. To
	plish this:
<u>PURC</u>	HASE THE FOLLOWING FROM YOUR PHARMACY:
One (1	l) 8.3oz bottle of miralax powder
Two (2	2) bottles of fleet water saline enemas (GREEN BOX)
PREP I	NSTRUCTIONS:
1.	Evening before procedure, Mix one 8.3oz bottle of miralax powder with 64oz of a clear
	liquid of your choice. Drink one 8oz glass every 15-20 minutes until you drank the entire
	64oz.
2.	NOTHING to eat after 12:00AM (mid-night). You may have clear liquids up until
	3HOURS before procedure time.
3.	On the day of procedure, TWO (2) Hours before leaving home, (or at
	am/pm), use the first fleet's enema. Hold it for as long as possible, and
	then expel the enema. After the first enema is completed, follow the same procedure
	using the 2 nd enema. After both enemas are completed, you may leave for your
	scheduled sigmoidoscopy procedure.
4.	You must report to our surgery center atam/pm.
<u>Impo</u>	<u>rtant:</u>
We M	UST have the referral for your sigmoidoscopy at least three (3) days before your
sigmoi	doscopy procedure.
	If you have any questions please call us at:
	215-402-0600

This is a procedure in which a flexible fiberoptic tube is placed in the rectum to view the rectum and bowel. The tube bends as the bowel bends, the doctor is able to look for causes of bleeding or discomfort, such as hemorrhoids, diverticulosis, colitis, polyps or cancer.

The sigmoidoscopy is done while you are lying on your side. The sigmoidoscopy usually takes from 5 to 15 minutes. Biopsy reports are usually available in one week to 10 days.

Revised 11/30/2022