

USDH PREPROCEDURE MEDICATION INSTRUCTIONS

NAME: _____ MR#/DOB _____

For your safety, your procedure may be rescheduled if you do not follow these directions

Anticoagulant/Blood Thinners: No need to stop prescribed aspirin or NSAIDs

STOP 2 DAYS BEFORE
Xarelto® (Rivaroxaban)
Eliquis® (Apixaban)
Pradaxa® (Dabigatran)
Savaysa® or Lixiana® (Edoxaban)

STOP 5 DAYS BEFORE
Plavix® (Clopidogrel)
Effient® (Prasugrel)
Brilinta® (Ticagrelor)
Coumadin® (Warfarin)

Other _____

GLP-1 Agonists OR GLP-1 Agonist/GIP combo – diabetes/weight loss/heart medications

STOP 1 DAY BEFORE
Victoza® and Saxenda® (Liraglutide)
Byetta® (Exenatide)

STOP 7 DAYS BEFORE
Ozempic®, Wegovy®, Rybelsus® (Semaglutide)
Bydureon® (Exenatide)
Trulicity® (Dulaglutide)
Mounjaro®, Zepbound® (Tirzepatide)

COLONOSCOPY PATIENTS ONLY :

Contact your provider who manages your diabetes to discuss whether you need additional management

Do not take for 3 days before colonoscopy and day of colonoscopy

Invokana® (Canagliflozin)	Jardiance® (Empagliflozin)
Invokamet® (Canagliflozin/metformin)	Synjardy® (Empagliflozin/metformin)
Invokamet® XR (Canagliflozin/metformin XR)	Synjardy® XR (Empagliflozin/metformin XR)
Farxiga® (Dapagliflozin)	Glyxambi® (Empagliflozin/linagliptin)
Xigduo® (Dapagliflozin/metformin)	Trijardy® XR (Empagliflozin/linagliptin/metformin XR)
Xigduo® XR (Dapagliflozin/metformin XR)	Steglatro® (Ertugliflozin)
Qtern® (Dapagliflozin/saxagliptin)	Segluromet® (Ertugliflozin/metformin)
Qternmet® XR (Dapagliflozin/saxagliptin/metformin)	Steglujan® (Ertugliflozin/sitagliptin)

ALL DIABETIC PATIENTS:

____ For Insulin - take ½ the usual dose the day before the procedure and none the day of procedure.

____ For diabetes medications besides any listed above - do not take the day of the procedure.

OTHER MED INSTRUCTIONS: _____

***Take all other medications (except water or fluid pills) on the day of the procedure as you normally would by 3 hours prior to arrival time**

PROVIDER: _____