



Allegheny Regional Endoscopy
810 Valley View Blvd.
Altoona, PA 16602
Front desk: 814-946-5469 Fax: 814-946-4970

Your colonoscopy is scheduled on:

Clenpiq Prep Instructions

- Avoid iron supplements for 5 days prior to your procedure.
- Avoid eating whole-grain bread/cereals with nuts/seeds and corn for 5 days prior to your procedure.

Stage 1

- Consume *clear liquids only* (see list below) the day before your procedure
- At 3:00pm the day before your procedure, take 2 Dulcolax tablets (5mg each)
- At 5:00pm the day before your procedure, drink 1 entire bottle of Clenpiq prep. Over the next 5 hours, drink 5 additional 8-ounce glasses of clear liquids, at your own pace.
- Once Stage 1 is complete, continue to consume *clear liquids* until it is time for stage 2 of the prep.

Stage 2

- *7 hours before your scheduled procedure time*, drink the second entire bottle of Clenpiq prep. Over the next 2 hours, drink 3 additional 8-ounce glasses of clear liquids. *Be sure stage 2 is completed entirely within 2 hours.*
- Once both stages of the prep are complete, you may continue to consume *clear liquids until 2 hours before your scheduled procedure time.*

Notes

- It may take a few hours for the prep to start working.
- It is normal to feel very full or bloated during your prep. *If you experience nausea and/or vomiting, wait 30 minutes before resuming prep/liquids. If vomiting does not disappear once you begin to move your bowels, please call us immediately for further instructions.*

ACCEPTABLE CLEAR LIQUIDS

Water Gatorade Jell-O Kool-Aid Broth Popsicles
Lemonade Apple juice White grape juice White cranberry juice Soda (all)
Black coffee (no added cream or sugar) Iced tea Hot tea (no added sugar, honey)

****Do not consume any liquids which are red or purple in color****

Colonoscopy and Anesthesia

It is especially important that you read this information in its entirety and call to speak with us if you are unsure of any instructions. **Failure to comply with the below guidelines could result in the delay and/or cancellation of your procedures without question.**

- **YOU MUST** have a driver to sign you out of recovery after anesthesia. **You cannot use UBER, LYFT, Healthride or MediVan unless a responsible party is riding beside you.**
- **You may have CLEAR LIQUIDS up to 2 hours before your designated arrival time at our facility and no later.**
- Please refer to the prep instructions on the front of this sheet. **DO NOT follow pharmacy instructions.**
- **No alcoholic beverages are permitted during colon prep OR after midnight on the night before procedures.**

CHEWING TOBACCO PRODUCTS (INCLUDING ALL POUCHES), SMOKING, VAPING AND INHALATION OF MARIJUANA (medical and recreational) **ARE PROHIBITED** after midnight, on the night before your procedures. You should also avoid hard candy and chewing gum after midnight.

- **All oral piercings** must be removed before all procedures as well as any other jewelry. **NO EXCEPTIONS!**
- **All stimulators** (bladder, spinal etc.) should be turned “off” or placed in “surgery mode” if you are having a colonoscopy.
- Please refer to the personalized medication instructions provided to you. ***It is especially important to take medications for blood pressure and heart function the day of your procedures.***
- ***If you experience any of the following between now and the day of your procedure, please notify us immediately:***
 - 1. A change in health status***
 - 2. an ER visit/hospital admission***
 - 3. Testing or office visit for your heart, lungs, brain***

Please call 814-946-5469 Option 7 to report these changes, or if you have any additional questions pertaining to your procedure.

!! PLEASE SEE THE CHART BELOW FOR A LIST OF MEDICATIONS THAT REQUIRE A MANDATORY HOLD PRIOR TO PROCEDURES (see next page)!!

HOLD MEDICATION DAY OF PROCEDURE:

Carafate (Sulcrafate)

HOLD MEDICATIONS FOR 1 DAY PRIOR TO PROCEDURE:

Byetta (Exenatide)
Soliqua (Lixisenatide + Insulin)
Victoza/Saxenda (Liraglutide)

HOLD MEDICATIONS FOR 3 DAYS PRIOR TO PROCEDURE:

Brenzavvy (Bexagliflozin)
Farxiga (Depagliflozin)
Glyxambi (Empagliflozin/Linagliptin)
Inpefa (Sotagliflozin)
Invokana (Canagliflozin)
Invokamet (Canagliflozin/Metformin)
Invokamet XR (Canagliflozin/Metformin XR)
Jardiance (Empagliflozin)
Qtern (Dapagliflozin/Saxagliptin)
Qternmet XR (Dapagliflozin/Saxagliptin/Metformin XR)
Synjardy (Empagliflozin/Metformin)
Synjardy XR (Empagliflozin/Metformin XR)
Steglatro (Ertugliflozin)
Steglujan (Ertugliflozin/Sitagliptin)
Segluromet (Ertugliflozin/Metformin)
Tijardy XR (Empagliflozin/Linagliptin/Metformin XR)
Xigduo (Dapagliflozin/Metformin)
Xigduo XR (Dapagliflozin/Metformin XR)

HOLD MEDICATIONS FOR 7 DAYS PRIOR TO PROCEDURE:

Adipex-P/Lomaira (Phentermine)
Adlyxin (Lixisenatide)
Bydureon (Exenatide)
Mounjaro/Zepbound (Tirzepatide)
Ozempic (Semaglutide)
Qysmia (Phentermine and Topiramate)
Rybelsus
Trulicity (Dulaglutide)
Wegovy