



Allegheny Regional Endoscopy
810 Valley View Blvd.
Altoona, PA 16602
Front desk: 814-946-5469 Fax: 814-946-4970

Your colonoscopy is scheduled on:

Suflave Prep Instructions

- Avoid iron supplements for 5 days prior to your procedure.
- Avoid eating whole-grain bread/cereals with nuts/seeds and corn for 5 days prior to your procedure.

Stage 1

- Consume *clear liquids only* (see list below) the day before your procedure
- Open 1 flavor-enhancing packet of prep and pour into the provided bottle. Fill the bottle to the "fill line" with lukewarm water. Cap the bottle, shake until completely mixed and *refrigerate for up to 24 hours. DO NOT FREEZE*. You can also mix and use immediately if desired.
- In the early evening on the day before your procedure, drink 8 ounces of the prep solution every 15 minutes until it is gone. When finished, drink an additional 16 ounces of water.
- Once Stage 1 is complete, continue to consume *clear liquids* until it is time for stage 2 of the prep.
- *There should be a minimum of 4 hours between stage 1 and stage 2.*

Stage 2

- *8 hours before your scheduled procedure time*, open the second flavor-enhancing packet of prep and again pour into the provided bottle.
- Repeat the steps as in Stage 1 above but *be sure to complete this entire stage within 3 hours*.
- Once both stages of the prep are complete, you may continue to consume *clear liquids until 2 hours before your scheduled procedure time*.

Notes

- It may take a few hours for the prep to start working.
- It is normal to feel very full or bloated during your prep. *If you experience nausea and/or vomiting, wait 30 minutes before resuming prep/liquids. If vomiting does not disappear once you begin to move your bowels, please call us immediately for further instructions.*

ACCEPTABLE CLEAR LIQUIDS

Water	Gatorade	Jell-O	Kool-Aid	Broth	Popsicles
Lemonade	Apple juice	White grape juice	White cranberry juice	Soda (all)	
Black coffee (no added cream or sugar)		Iced tea	Hot tea (no added sugar, honey)		

Do not consume any liquids which are red or purple in color

Colonoscopy and Anesthesia

It is especially important that you read this information in its entirety and call to speak with us if you are unsure of any instructions. **Failure to comply with the below guidelines could result in the delay and/or cancellation of your procedures without question.**

- **YOU MUST** have a driver to sign you out of recovery after anesthesia. **You cannot use UBER, LYFT, Healthride or MediVan unless a responsible party is riding beside you.**
- **You may have CLEAR LIQUIDS up to 2 hours before your designated arrival time at our facility and no later.**
- Please refer to the prep instructions on the front of this sheet. **DO NOT follow pharmacy instructions.**
- **No alcoholic beverages are permitted during colon prep OR after midnight on the night before procedures.**

CHEWING TOBACCO PRODUCTS (INCLUDING ALL POUCHES), SMOKING, VAPING AND INHALATION OF MARIJUANA (medical and recreational) **ARE PROHIBITED** after midnight, on the night before your procedures. You should also avoid hard candy and chewing gum after midnight.

- **All oral piercings** must be removed before all procedures as well as any other jewelry. **NO EXCEPTIONS!**
- **All stimulators** (bladder, spinal etc.) should be turned “off” or placed in “surgery mode” if you are having a colonoscopy.
- Please refer to the personalized medication instructions provided to you. ***It is especially important to take medications for blood pressure and heart function the day of your procedures.***
- ***If you experience any of the following between now and the day of your procedure, please notify us immediately:***
 - 1. A change in health status***
 - 2. an ER visit/hospital admission***
 - 3. Testing or office visit for your heart, lungs, brain***

Please call 814-946-5469 Option 7 to report these changes, or if you have any additional questions pertaining to your procedure.

!! PLEASE SEE THE CHART BELOW FOR A LIST OF MEDICATIONS THAT REQUIRE A MANDATORY HOLD PRIOR TO PROCEDURES (see next page)!!

HOLD MEDICATION DAY OF PROCEDURE:

Carafate (Sulcrafate)

HOLD MEDICATIONS FOR 1 DAY PRIOR TO PROCEDURE:

Byetta (Exenatide)
Soliqua (Lixisenatide + Insulin)
Victoza/Saxenda (Liraglutide)

HOLD MEDICATIONS FOR 3 DAYS PRIOR TO PROCEDURE:

Brenzavvy (Bexagliflozin)
Farxiga (Depagliflozin)
Glyxambi (Empagliflozin/Linagliptin)
Inpefa (Sotagliflozin)
Invokana (Canagliflozin)
Invokamet (Canagliflozin/Metformin)
Invokamet XR (Canagliflozin/Metformin XR)
Jardiance (Empagliflozin)
Qtern (Dapagliflozin/Saxagliptin)
Qternmet XR (Dapagliflozin/Saxagliptin/Metformin XR)
Synjardy (Empagliflozin/Metformin)
Synjardy XR (Empagliflozin/Metformin XR)
Steglatro (Ertugliflozin)
Steglujan (Ertugliflozin/Sitagliptin)
Segluromet (Ertugliflozin/Metformin)
Tijardy XR (Empagliflozin/Linagliptin/Metformin XR)
Xigduo (Dapagliflozin/Metformin)
Xigduo XR (Dapagliflozin/Metformin XR)

HOLD MEDICATIONS FOR 7 DAYS PRIOR TO PROCEDURE:

Adipex-P/Lomaira (Phentermine)
Adlyxin (Lixisenatide)
Bydureon (Exenatide)
Mounjaro/Zepbound (Tirzepatide)
Ozempic (Semaglutide)
Qysmia (Phentermine and Topiramate)
Rybelsus
Trulicity (Dulaglutide)
Wegovy